



ISSN: 0975-833X

Available online at <http://www.journalcra.com>

International Journal of Current Research  
Vol. 11, Issue, 03, pp.2167-2170, March, 2019

DOI: <https://doi.org/10.24941/ijcr.34772.03.2019>

INTERNATIONAL JOURNAL  
OF CURRENT RESEARCH

## RESEARCH ARTICLE

### DEPRESSION, ANXIETY AND STRESS AMONG HIV PATIENTS IN EAST COAST MALAYSIA

<sup>1,\*</sup>Azwanis Binti Abdul Hadi, <sup>1</sup>Nur Syuhada Binti Mohd Mustapha, <sup>1</sup>Nur Farhana Binti Ahmad Hafidzi, <sup>1</sup>Nur Sahira Binti Shaari, <sup>2</sup>Hafizah Binti Pasi, <sup>3</sup>Iskandar Firzada Osman, <sup>4</sup>Mohd Daud Bin Che Yusof

<sup>1</sup>Department of Family Medicine, International Islamic University Malaysia  
<sup>2</sup>Department of Community Medicine, International Islamic University Malaysia  
<sup>3</sup>Klinik Kesihatan Jaya Gading, Kuantan, Pahang, Malaysia  
<sup>4</sup>Klinik Kesihatan Beserah, Kuantan, Pahang, Malaysia

#### ARTICLE INFO

##### Article History:

Received 25<sup>th</sup> December, 2018  
Received in revised form  
14<sup>th</sup> January, 2019  
Accepted 18<sup>th</sup> February, 2019  
Published online 31<sup>st</sup> March, 2019

##### Key Words:

HIV, Depression, Anxiety, Stress,  
Community, Malaysia.

##### \*Corresponding author:

Azwanis Binti Abdul Hadi

Copyright © 2019, Azwanis Binti Abdul Hadi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Azwanis Binti Abdul Hadi, Nur Syuhada Binti Mohd Mustapha, Nur Farhana Binti Ahmad Hafidzi, et al. 2019. "Depression, Anxiety and Stress among HIV Patients in East Coast Malaysia", *International Journal of Current Research*, 11, (03), 2167-2170.

#### ABSTRACT

**Introduction:** HIV infection is a chronic disease with multiple challenges that could affect the psychological well-being of patients. **Objective:** The objective of this research was to determine the prevalence of depression, anxiety and stress among HIV patients and its significant associated factors. **Methodology:** This is a cross-sectional study which was carried out in two health centers in a town in the East Coast of peninsular Malaysia. A total of 42 patients with HIV were recruited via convenience sampling. Those who fulfilled the inclusion and exclusion criteria completed a validated set of Malay version of DASS-21 questionnaire. Data was analyzed using IBM SPSS Statistics. **Results:** The prevalence of depression, anxiety and stress among adult HIV patients in community follow-up were 33.3%, 28.6% and 21.4% respectively. Marital status was significantly associated with both depression and stress. While CD4 count was found to be significantly associated with anxiety.

## INTRODUCTION

Patients who are diagnosed with Human immunodeficiency virus (HIV) could be affected psychologically. HIV is a lifelong disease that necessitates a patient to be compliant with multiple medications. It is also a disease well known for its social stigma. However, there are many other reasons that may lead to psychological disorder such as the adverse effects of antiretroviral drugs (Dubé *et al.*, 2005) or the serostatus of the patient (Kelly *et al.*, 1998). The most common psychological disorder that are usually under diagnosed is depression. As a result, the repercussions of mental illness on HIV patients may raise several issues such as HAART compliance, defaulted follow up and deteriorating quality of life that eventually leads to increase in mortality rate (Simoni *et al.*, 2011). In Malaysia, with the initiation of comprehensive treatment cascade such as the screening process, Human Anti-Retroviral Therapy (HAART) treatment and continuous follow up for viral load suppression with CD4 count had achieve great achievement in reducing half the case of HIV infected patient (MOH, 2016). This is evidenced by 22 cases per 100,000 population in 2010 reducing to 11.2 cases per 100,000 population in 2014 (MOH, 2016).

However, there are still approximately 3517 of Malaysians infected with HIV In the year 2014, and there might be a significant number of them who are inflicted with a degree of psychological illness. According to the Malaysian National Health and Morbidity Survey, the prevalence of psychological illness was 11% in a normal population. Despite the worrying prevalence of this psychological illness, it is often missed in the primary care setting leading to difficulties for physicians to provide an optimal management of the disease to their patients (Simoni, 2011).

## MATERIALS AND METHODS

**Ethical consideration:** Ethical approval was obtained from Ethical Committee of Kuliyyah of Medicine, International Islamic University Malaysia [IREC 2017-010] and Medical Research and Ethics Committee, Ministry of Health Malaysia [NMRR-17-746-35329].

**Study design, population and sampling method:** A cross-sectional study were conducted in two health clinics in Kuantan, Malaysia. We included adult HIV patients that were on highly active anti-retroviral therapy (HAART) and who understood Malay.

The study was conducted over a period of 6 weeks from 17<sup>th</sup> July 2017 to 27<sup>th</sup> August 2017. Patients were recruited using convenience sampling.

**Study tool:** This study used a self-administered questionnaire which consisted of three parts. Part A: socio-demographic information of the patient, Part B: Malay version of the Depression, Anxiety and Stress Scale 21-item (DASS-21), Part C: related medical records of the patient.

Patient's informed consent form were attached together with the questionnaire. Depression, Anxiety and Stress Scale 21-item (DASS-21) is a self-reporting tool which measures three related negative emotional states, which are depression, anxiety and stress. Each components has a different scoring and cut-off value. Among the advantages of DASS-21 Bahasa Melayu version is that it can be administered to both clinical and non-clinical subjects while also possessing good psychometric properties for clinical subject (Musa, 2009). This questionnaire is simple and easy to be administered and has been widely used in Malaysia (Musa, 2012).

**Statistical Analysis:** The statistical analysis was performed using software IBM SPSS Statistics for Windows, Version 23.0. Descriptive statistics were performed for all demographic factors and related medical records. While bivariate analysis was done to study the relations between variables.

## RESULTS

A total of 42 patients with HIV were recruited in this study. Majority of the respondents were male, Malays, married, living with partner/family, with secondary school level of education and belonging to the low monthly household income group (< RM 1000 per month). The mean age for the respondents were 40 years old. Majority of the respondents' had CD4 count more than 350 cells per mm<sup>3</sup> with undetectable viral load. Majority have other medical co-morbidities and are on less than 5 types of medications. Table 1 shows the characteristics of the HIV patients.

**Prevalence of depression, anxiety and stress:** The prevalence of depression, anxiety and stress among adult HIV patients on HAART were 33.3%, 28.6% and 21.4% respectively. From this, less than 10% were found to be from the extremely severe subscale for each group (7.1%, 9.5% and 2.4% respectively).

**Relationship of sociodemographic characteristics and related medical records of the respondents with depression, anxiety and stress level:** Patients' data were analysed using the chi-square test to determine the relationship of sociodemographic characteristics and related medical records of the respondents with depression, anxiety and stress levels as shown in Table 2, 3, 4. Marital status was significantly associated with both depression and stress. While CD4 count was found to be significantly associated with anxiety.

## DISCUSSION

The prevalence of depression in this study was 33.3% which was comparable with studies done in another part of Malaysia

**Table 1. Socio-demographic characteristics and related medical records of the respondents (n=42)**

Variables	N (%)	
<b>Age (years)</b>	40.62	(11.00)*
<b>Gender</b>		
Male	30	(71.4)
Female	12	(28.6)
<b>Marital status</b>		
Married	20	(47.6)
Single	13	(31)
Widow/widower	9	(21.4)
<b>Living arrangement</b>		
Alone	5	(11.9)
With partner/family	36	(85.7)
With friends	1	(2.4)
<b>Religion</b>		
Islam	40	(95.2)
Buddhist	2	(4.8)
<b>Ethnicity</b>		
Malay	40	(95.2)
Chinese	2	(4.8)
<b>Highest education level</b>		
Primary	4	(9.5)
Secondary	21	(50.0)
Tertiary	17	(40.5)
<b>Occupation</b>		
Professional	12	(28.6)
Supporting staff	11	(26.2)
Part-time worker	6	(14.3)
Housewife	5	(11.9)
Retiree	1	(2.4)
Student	1	(2.4)
Not working	6	(14.3)
<b>Monthly household income</b>		
≤ RM 1000	21	(50.0)
RM 1001- RM 1500	3	(7.1)
RM 1501- RM 2000	4	(9.5)
RM 2001- RM 2500	4	(9.5)
RM 2501- RM 3000	1	(2.4)
≥RM 3000	9	(21.4)
<b>CD4 count (cells per mm<sup>3</sup>)</b>		
≤ 350	18	(42.9)
> 350	24	(57.1)
<b>Viral load</b>		
Undetectable	24	(57.1)
Detectable	14	(33.3)
<b>Comorbidities</b>		
No	14	(33.3)
Yes	28	(66.7)
<b>Number of medications</b>		
< 5	29	(69.0)
5-10	13	(31.0)

**Table 2. Relationship of socio-demographic characteristics and related medical records of the respondents with depression level**

Variables	Yes		No		P-value
	n	%	n	%	
<b>Marital status</b>					<b>0.008</b>
Married	7	(35.0)	13	(65.0)	
Single	7	(53.8)	6	(46.2)	
Widower	0	(0)	9	(100)	

**Table 3: Relationship of socio-demographic characteristics and related medical records of the respondents with anxiety level**

Variables	Yes		No		P-value
	n	%	n	%	
<b>CD4 count (cells per mm<sup>3</sup>)</b>					0.048
≤ 350	9	(50.0)	9	(83.3)	
>350	4	(16.7)	20		

**Table 4. Relationship of socio-demographic characteristics and related medical records of the respondents with stress level**

Variables	Yes		No		P-value
	n	%	n	%	
<b>Marital status</b>					<b>0.007</b>
Married	2	(20.0)	18	(90.0)	
Single	6	(46.2)	7	(53.8)	
Widower	0	(0)	9	(21.4)	

and Korea (Radzniwan *et al.*, 2016; Kee *et al.*, 2015). Interestingly, it was lower than the studies in China, South Africa, Australia and Sudan in which their prevalence of depression was in the range of 41% to 63% (Li *et al.*, 2016; Elbadawi, 2017; Heywood, 2016; Freeman *et al.*, 2014). The differences could be because of the different tools used and the different factors in their population. When we compared head-to-head with a study done in the northern part of this country, we found that the prevalence of depression and stress in our study (33.3%, 21.4%) is similar with theirs (36.9%, 26.7%). However, the prevalence of anxiety is much higher (45.1%) in their study as compared to ours (28.6%) (Radzniwan *et al.*, 2016). The prevalence of their participants in the extremely severe group for depression, anxiety and stress were also higher as compared to our study. While only less than 10% of our patients were in the extremely severe group for depression, anxiety and stress; they reported extremely severe depression and anxiety as high as 14.5% and 26.9% respectively (Radzniwan *et al.*, 2016).

The main difference between the studies is that their participants were patients under follow-up at the hospital while our participants were under community health clinics follow-up. Usually the patients treated in the community are less ill and have more straight-forward treatment. In our study, being single is significantly associated with both depression and stress. This concurred with a study in Iran which also found that depression is higher in the single (married, widowed and divorced) women (Saadat *et al.*, 2015). This could be because of the poor support that patients receive. In a study done in Australia, it was found that social support was a protective factor. And among all the supports that they studied, emotional support was the most significant in reducing psychological problems (Heywood, 2016). So, emotional support from partners or spouse acts as a buffer against the stress that being a patient with HIV can bring (Saadat *et al.*, 2015). But interestingly in Korea and Sudan, they found that their patients who were married or currently living with someone were more likely to have anxiety and depressive symptoms than those who were single. It was found that patients who were infected by their spouse or partners are highly affected psychologically as well as decreased sexual desire (Kee *et al.*, 2015; Saadat *et al.*, 2015). This study found a significant relationship between CD4 count with anxiety level. Patients with higher CD4 count (>350) had lower anxiety level as compared with patients with lower CD4 count. This corresponds with a research in Australia where they also reported significant association between CD4 count and anxiety. CD4 count is an important laboratory indicator of a person's immune function and is a predictor for HIV disease progression. A lower CD4 count correlate with more HIV associated illnesses and poorer quality of life.

## Conclusion

There are presence of depression, anxiety and stress among HIV patients on community follow-up. There are even a small

but important percentage of them who were suffering from extremely severe depression.

We found that marital status and CD4 count were significantly associated with depression, anxiety and stress. Hence, early detection of the disorders and providing proper psychiatric management for HIV patients is important to ensure the quality of life of these patients.

**Limitations:** This study has limitation on the small sample size and design as it was a cross-sectional study and the selection of participants were based on convenience sampling. There is a possibility of response bias using a guided questionnaire.

**Acknowledgement:** The authors would like to acknowledge International Islamic University Malaysia for supporting this study.

**Funding declaration:** No Funding

## REFERENCES

- Dubé, B., Benton, T., Cruess, D. G., & Evans, D. L. (2005). Neuropsychiatric manifestations of HIV infection and AIDS. *J Psychiatry Neurosci* 2005, 30(4), 237-246.
- Elbadawi, A., & Mirghani, H. 2017. Depression among HIV/AIDS Sudanese patients: a cross-sectional analytic study. *Pan Afr Med J*, 26, 43. doi:10.11604/pamj.2017.26.43.10919
- Freeman, M., Nkomo, N., Kafaar, Z., & Kelly, K. 2014. Mental disorder in people living with HIV/Aids in South Africa. *South African Journal Of Psychology*, 38(3), 489-500.
- Heywood, W., & Lyons, A. 2016. HIV and Elevated Mental Health Problems: Diagnostic, Treatment, and Risk Patterns for Symptoms of Depression, Anxiety, and Stress in a National Community-Based Cohort of Gay Men Living with HIV. *AIDS Behav*, 20(8), 1632-1645. doi:10.1007/s10461-016-1324-y
- Kee, M. K., Lee, S. Y., Kim, N. Y., Lee, J. S., Kim, J. M., Choi, J. Y., Kim, S. S. 2015. Anxiety and depressive symptoms among patients infected with human immunodeficiency virus in South Korea. *AIDS Care*, 27(9), 1174-1182. doi:10.1080/09540121.2015.1035861
- Kelly, B., Raphael, B., Judd, F., Perdices, M., Kernutt, G., Burrow, G. D., Dunne, M. (1998). Psychiatric disorder in HIV infection. *Australian and New Zealand Journal of Psychiatry*, 32, 441-453.
- Li, J., Mo, P. K., Kahler, C. W., Lau, J. T., Du, M., Dai, Y., & Shen, H. 2016. Prevalence and associated factors of depressive and anxiety symptoms among HIV-infected men who have sex with men in China. *AIDS Care*, 28(4), 465-470. doi:10.1080/09540121.2015.1118430
- MOH. 2016. Malaysia GARPR 2016.
- Musa, R. 2012. *Profile of Malaysian version of the Depressive, Anxiety and Stress Scale 42-item (DASS-42)* (Vol. 20).
- Musa, R., M. A, S., & Ain M, N. 2009. *Validation and Psychometric Properties of Bahasa Malaysia Version of the Depression Anxiety And Stress Scales (DASS) Among Diabetic Patients* (Vol. 18).
- Radzniwan, R., Alyani, M., Aida, J., Khairani, O., Nik Jaafar, N. R., & Tohid, H. 2016. Psychological status and its clinical determinants among people living with HIV/AIDS (PLWHA) in Northern Peninsular Malaysia.

- HIV & AIDS Review*, 15(4), 141-146. doi:10.1016/j.hivar.2016.11.002
- Saadat, M., Behboodi, Z. M., & Saadat, E. 2015. Comparison of depression, anxiety, stress, and related factors among women and men with human immunodeficiency virus infection. *J Hum Reprod Sci*, 8(1), 48-51. doi:10.4103/0974-1208.153128
- Simoni, J. M., Safren, S. A., Manhart, L. E., Lyda, K., Grossman, C. I., Rao, D., Wilson, I. B. 2011. Challenges in addressing depression in HIV research: assessment, cultural context, and methods. *AIDS Behav*, 15(2), 376-388. doi:10.1007/s10461-010-9836-3

\*\*\*\*\*