



RESEARCH ARTICLE

PREVALENCE OF LOW BACK PAIN AMONG HEALTH SCIENCES STUDENTS IN SRINAGAR: A CROSS SECTIONAL STUDY

^{1,*}Gazalla Altaf, ²Mehrajud Din Bhat and ¹Mehak Anand

¹Department of Pedodontics and Preventive Dentistry, Sudha Rustagi College of Dental Sciences and Research, Faridabad, Haryana, India

²Department of Prosthodontics, Sudha Rustagi College of Dental Sciences and Research, Faridabad, Haryana, India

ARTICLE INFO

Article History:

Received 18th April, 2019

Received in revised form

14th May, 2019

Accepted 12th June, 2019

Published online 31st July, 2019

Key Words:

Health sciences
Students.

*Corresponding author: Gazalla Altaf

Copyright © 2019, Gazalla Altaf et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Gazalla Altaf, Mehrajud Din Bhat and Mehak Anand, 2019. "Prevalence of low back pain among health sciences students in Srinagar: A cross sectional Study", *International Journal of Current Research*, 11, (07), 5674-5677.

ABSTRACT

Aim: To determine the prevalence of low back pain (LBP) among health sciences students and to identify the associated factors. **Methods:** Cross-sectional study was conducted among 1163 students from five health sciences colleges during the academic year 2016–2017. Self-administered questionnaire was conducted and included 4 sections: demographic characteristics, risk factors, Nordic musculoskeletal questionnaire and Oswestry disability questionnaire. Data were analyzed using SPSS. **Results:** Mean age was 20.74 ± (1.59 years). 70.9% of students were female. Lifetime prevalence of LBP was 56.6%, 12-month prevalence 48.8%, and point prevalence 21.2%. Dentistry students had highest lifetime prevalence of LBP (67.6%) with significant p value (<0.001). Male were found to have higher lifetime prevalence compared to female (p ≤ 0.001). Spending more than 10 h on computer or tablet was significantly associated with LBP (OR 2.19; 95% CI 1.30–3.70; p = 0.003). Feeling discomfort on bed was associated with LBP (OR 1.81; 95% CI 1.38–2.38; p ≤ 0.001). Uncomfortable college furniture was associated with LBP (OR 1.40; 95% CI 1.09–1.79; p = 0.008). Using heavy backpack was found to be associated with LBP (OR 1.49; 95% CI 1.01–2.03; p = 0.011). Most of students LBP (90.3%) found to cause minimal disability on Oswestry scale. **Conclusion:** Dentistry students at higher risk of developing LBP compared to other students

INTRODUCTION

Job-related musculoskeletal disorders (MSDs) usually happen over a period of time, resulting from repeated work-load exposures. The neck, low back, and upper limbs are commonly vulnerable to MSDs (Bernacki et al., 1999). The life time prevalence of LBP is reported to be high, affecting nearly 80% of people at some time in their adult life, and the point prevalence is ranging from 30% to 50% (Jin et al., 2004). For example, in France more than half of the French population experienced LBP at least one day in the previous 12 months (Gourmelen et al., 2007). Low back pain is commonly recognized "non-specific" as no organic source was found to cause the pain (Urquhart et al., 2011). Frequent initial onset of back pain appears in age of 30 years, and the peak is between 45 and 60 years (Anand et al., 2013). Health science students are vulnerable to stress and prolonged time of studying and training which will make them predisposed for having LBP. Reported overall prevalence of LBP in health science students ranges from 40.1 to 57.9%, but there is controversy about which college has highest prevalence of LBP (Alshagga et al., 2013). According to Nordic classification of LBP, lifetime prevalence, 1-year prevalence and point prevalence of LBP among: medical students were 73.4. 46.1–59.9 and 27.2%:

physical therapy students 69.2–82.3, 63.2–73.7 and 27.6%; and nursing students 79, 71 and 30%, respectively (Moroder et al., 2011). LBP might be chronic, and this will affect future health professionals in the provision of health care to patients. Studies have shown conflicting results on the effect of identifiable risk factors in health science students on LBP. Identified risk factors for LBP included gender, age, weight, general health status, socioeconomic status, smoking, year of study, psychosocial factors (feeling very sad, overwhelmed, exhausted), history of back pain, history of back trauma, family history of treated back pain, use of heavy backpack, physical fitness, prolonged sitting time, bad postural habits, short sleep hours, discomfort in bed and college furniture (Bernacki et al., 1999; Jin et al., 2004; Anand et al., 2013; Alshagga et al., 2013; Moroder et al., 2011). If risk factors are well recognized and understood, this would help in establishing preventive measures of LBP and modifying risk factors which will lead to better quality of life for future health professionals. No studies have been done in our region about the prevalence of low back pain among health science students. Further risk factors will be tested in this study. Aim of this

Quantitative cross-sectional study was conducted at 5 health science colleges (medicine, dentistry, pharmacy, nursing and college of applied medical sciences “physical therapy, occupational therapy, clinical laboratory and clinical nutrition”) at Srinagar, which was conducted during the academic year 2017–2018. The study included 1163 students and excluded students who have spinal deformities or had undergone back surgery. Self-administered questionnaire was distributed. Instruction was explained by one of the authors and the data collector who was well informed about all the parts of questionnaire. Completeness of questionnaires was checked by the authors. The questionnaire contains 52 questions in 3 pages divided into 4 sections: demographic characteristics such as age, gender, weight, height, college, year at college. Risk factors included smoking, physical activity, coffee drinking, hours spent on using computer, position while studying, sleeping hours, comfort of bed to back, comfort of college furniture to bed, using heavy backpack, being overwhelmed, feeling exhausted and feeling sad. LBP was assessed using Nordic musculoskeletal questionnaire. Disability was classified based on Oswestry questionnaire. Nordic and Oswestry questionnaires were previously tested for validity and reliability (Kuorinka *et al.*, 1987; Fairbank and Pynsent, 2000). Data were analyzed by using Statistical Package for Social Studies (SPSS 22; IBM Corp., New York, NY, USA). Continuous variables were expressed as mean \pm standard deviation, and categorical variables were expressed as percentages. The t-test was used for continuous variables, and Chi-square test was used for categorical variables. Odds ratio (OR) with their 95% confidence interval (CI) was used for assessing the risk factors for LBP using the univariate analysis. A p value <0.05 was considered statistically significant.

RESULTS

One thousand and fifty-two participants out of 1163 were enrolled in this study. Mean age of participants \pm (standard deviation) was $20.74 \pm (1.59)$ years. Majority of participants were female (70.9%). Demographic characteristics are shown in Table 1. Overall lifetime prevalence of LBP among health sciences undergraduate was 56.6%, 12-month prevalence 48.8%, and point prevalence 21.2%. Dentistry students had significantly highest lifetime, 12-month and point prevalence compared to other students. Males had significantly higher lifetime prevalence of LBP than females (Table 3). Risk factors associated significantly with point prevalence of LBP are hours spent on using computer or tablet ($p = 0.002$), sleeping on uncomfortable bed ($p < 0.001$), uncomfortable college furniture ($p = 0.005$), using heavy backpack ($p = 0.01$), feeling overwhelmed ($p < 0.001$), feeling sad ($p < 0.001$), previous history of back trauma ($p < 0.001$) and family history of low back pain treated by a doctor ($p < 0.001$).

DISCUSSION

The overall lifetime prevalence of LBP among undergraduates was found to be high (56.6%), which is within the range of previous studies (40.1–57.9%). Further, 1-year prevalence was reported to be 48.8% and point prevalence 21.2%, which is higher than the results reported in Hafeez *et al.*'s study (21%, 13.1%) (Nordin *et al.*, 2014).

Table 1. Demographic characteristics

Characteristic	n(%)
Gender	
Male	306 (29.1)
Female	747 (70.9)
Age	
≤ 19	261 (24.8)
20	302 (28.7)
21	171 (16.2)
≥ 22	318 (30.2)
Colleges and subspecialties	
Medicine	328 (31.1)
Dentistry	145 (13.8)
Pharmacy	217 (20.6)
Nursing	120 (11.4)
Physicaltherapy	123 (11.7)
Occupationaltherapy	24 (2.3)
Clinical nutrition	49 (4.7)
Clinical laboratory	47 (4.5)
Academic year	
First	281 (26.7)
Second	275 (26.1)
Third	180 (17.1)
Fourth	152 (14.4)
Fifth	160 (15.5)
Tobacco	
Yes	48 (4.6)
Ex-smoker	9 (0.9)
No	996 (94.6)
Physical activity	
Active	403 (38.3)
Inactive	648 (61.5)
BMI	
< 20	212 (20.1)
20–25	453 (43)
> 25	296 (28.1)
Hours spent on using computer or tablet (h)	
< 4	228 (21.7)
4–10	391 (37.1)
> 10	434 (41.2)
Overwhelmed	
Yes	447 (42.5)
No	590 (56)
Sadness	
Yes	403 (38.3)
No	648 (61.5)
Exhausted	
Yes	608 (57.7)
No	442 (42)

In our study, dentistry students showed significantly highest lifetime, 1-year and point prevalence of LBP compared to other students. This could be attributed to the nature of their clinical training. As dentists use prolonged sitting and standing during their job, apply awkward posture and repetitive movements, many loads are exerted to the lumbar spine. It is believed that the higher muscular demand may lead to fatigue and consequently increase the risk of LBP in dentists. In a systematic review, it was reported that the prevalence of general musculoskeletal pain ranges between 64% and 93% and the most prevalent regions for pain in dentists have been shown to be the back (36.3–60.1%) (Udoye and Aguwa, 2007). In our study we found that Males suffered more pain than females. These results are similar to the results of study done by Al Wazzan *et al.* with findings - 58.24% males and 41.75% females suffered with neck and lowered back pain (Al Wazzan *et al.*, 2001). Also a study by Aasa *et al.* demonstrated that males reported a higher prevalence of LBP than women (Aasa *et al.*, 2005). Though a study by Leijon Mulder *et al.* concluded that female were more likely to report LBP than men (Leijon and Mulder, 2009). BMI was not significantly associated with LBP. which is consistent with previous studies

Table 2 Comparison of low back pain prevalence among different

	Medicine	Dentistry	Pharmacology	Nursing	Physical therapy
Lifetime prevalence	63.7	67.6	45.2	50	52
12-month prevalence	55.5	60.3	36.4	42.5	47.2

Coffee consumption increased in students having LBP, but in this study no significant correlation was found between coffee consumption and LBP. Most of students (61.4%) are physically inactive, and no significant association was found between physical activity and LBP. It is reported that prolonged sitting time using computer or tablet is strongly associated with LBP (Hafeez *et al.*, 2013). In this study, the result is consistent with the literature and it shows significantly higher point prevalence among students spending more than 10 h using computer or tablet compared with other groups. Prolonged sitting increases compression load on the spine. Position while studying and sleeping hours were not associated

with incidence of LBP. Hafeez and Yucel found an association between LBP and history of trauma of back, but it was not significant (Hafeez *et al.*, 2013). In contrast, Alshagga reported significant association between trauma and family history of musculoskeletal disorders with LBP¹⁷. In this study, previous history of trauma of back and family history of treated LBP were strongly associated with incidence of LBP ($p < 0.001$).

Conclusion

Health science students reported high prevalence of LBP. Dentistry students are at higher risk to develop LBP compared with other colleges. Lectures about preventive measures should

be included in health-related courses. Students should take a break during using computer or tablet and they should do stretching exercises to back. What we need is not preventing its incidence and recurrence by including preventive aerobic and relaxation exercises in weekly activities of dentists. This will help in improving the quality of work in their clinical practice.

REFERENCES

- Aasa U, Barnekow-Bergvist M, Angquist KA, Brulin C. 2005. Relationship between work-related factors and disorders in the neck-shoulder and low-back region among female and male ambulance personnel. *J Occup Health*, 47:481-489.
- Al Wazzan KA, Almas K, Al Shethri SE, Al-Qahtani MQ. 2001. Back & neck problems among dentists and dental auxiliaries. *J Contemp Dent Pract.*, 2:17-30
- Alshagga M, Nimer A, Yan L, Ibrahim I, Al-Ghamdi S, Radman Al-Dubai S. 2013. Prevalence and factors associated with neck, shoulder and low back pains among medical students in a Malaysian Medical College. *BMC Res Notes* 6(1):244
- Alshagga M, Nimer A, Yan L, Ibrahim I, Al-Ghamdi S, Radman Al-Dubai S. 2013. Prevalence and factors associated with neck, shoulder and low back pains among medical students in a Malaysian Medical College. *BMC Res Notes*, 6(1):244
- Anand T, Aggarwal N, Kishore J, Ingle G. 2013. Low back pain and associated risk factors among undergraduate students of a medical college in Delhi. *Educ Health* 26(2):103
- Bernacki EJ, Guidera JA, Schaefer JA. et al. 1999. An ergonomics program de-signed to reduce the incidence of upper extremity work related musculo-skeletal disorders. *J Occup Environ Med.*, 41: 1032–1041.
- Gourmelen J, Chastang JF, Oztguler A, Lanoë L, Ravaud JF, Leclerc A. 2007. Frequency of low back pain among men and women aged 30 to 64 years in France. Results of two national surveys. *Ann Readapt Med Phys.*, 50:640–644. doi:10.1016/j.annrmp.2007.05.00
- Hafeez K, Ahmed Memon A, Jawaid M, Usman S, Usman S, Haroon S. 2013. Back pain—are health care undergraduates at risk? *Iran J Public Health* 42(8):819–825
- Hulya Yucel, Perihan Torun, 2016. Incidence and risk factors of low back pain in students studying at a health university. *Bezmialem Sci.*, 4(1):12–18
- Jin K Vitae, Sorock, Courtney TK. 2004. Prevalence of low back pain in three occupational groups in Shanghai, People's Republic of China. *J Safety Res.*, 35:23-28. doi: 10.1016/j.jsr.2003.11.002
- Kuorinka I, Jonsson B, Kilbom A, Vinterberg H, Biering-Sørensen F, Andersson G. et al. 1987. Standardised Nordic questionnaires for the analysis of musculoskeletal symptoms. *Appl Ergon.*, 18(3):233–237
- Leijon O, Mulder M. 2009. Prevalence of low back pain and concurrent psychological distress over a 16-year period. *Occup Environ Med.*, 66:137-139. doi: 10.1136/oem.2008.04033
- Moroder P, Runer A, Resch H, Tauber M. 2011. Low back pain among medical students. *Acta Orthop Belg.*, 77:88–92
- Nordin NAM, Singh DKA, Kanglun L. 2014. Low back pain and associated risk factors among health science undergraduates. *Sains Malays*, 43(3):423–428
- Udoye CI, Aguwa EN. 2007. Musculoskeletal symptoms: a survey amongst a selected Nigerian dentists. *Int J Dent Sci.*, 5:1-5
- Urquhart D, Berry P, Wluka A, Strauss B, Wang Y, Proietto J. et al. 2011. Young investigator award winner. *Spine* 36(16):1320–1325
