



ISSN: 0975-833X

Available online at <http://www.journalcra.com>

INTERNATIONAL JOURNAL
OF CURRENT RESEARCH

International Journal of Current Research
Vol. 11, Issue, 08, pp.6115-6118, August, 2019

DOI: <https://doi.org/10.24941/ijcr.35801.08.2019>

RESEARCH ARTICLE

THE IMPACT OF ART THERAPY IN EVOLUTION OF PARASITIC DIARRHOEA IN SEROPOSITIVE PATIENT

**Dr. Rakesh Kumar (Associate Professor), Dr. Shailesh Kumar (Additional Professor),
Dr. Somya Sinha, Dr. Sweta Muni (Assistant Professor), Dr. Namrata Kumari (Professor) and
Prof (Dr.) Shahi, S.K.**

IGIMS, Patna, India

ARTICLE INFO

Article History:

Received 18th May, 2019
Received in revised form
16th June, 2019
Accepted 25th July, 2019
Published online 31st August, 2019

Key Words:

Essentials, Peace Culture, Faculty Encounter,
State Universities, Philippines

*Corresponding author: *Somya Sinha*

Copyright © 2019, *Rakesh Kumar et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: *Dr. Rakesh Kumar (Associate Professor), Dr. Shailesh Kumar (Additional Professor), Dr. Somya Sinha, et al. 2018.* "The impact of art therapy in evolution of parasitic diarrhoea in seropositive patient", *International Journal of Current Research*, 11, (08), 6115-6118.

ABSTRACT

Human immunodeficiency virus (HIV) infection has altered both the epidemiology and outcome of enteric opportunistic parasitic infections. This study was done to determine the prevalence and species/genotypes of intestinal coccidian and microsporidial infections among HIV/AIDS patients with diarrhea and/or a history of diarrhea alternately with an asymptomatic interval, and their association with ART therapy. This cross sectional study was done from May 2015 to March 2018 in IGIMS, Patna. Sociodemographic data and a history of diarrhea were collected by interviewing 234 HIV patients (146 males and 94 females). Using routine microscopy, trichrome staining, modified ZN staining incidence of coccidian parasites were found to be as *Cryptosporidium* 21.8% , followed by *Strongyloides* 17.6%, microsporidia, *Isospora*, *Giardia*, *Trichuris*, and *taenia*. ART treatment was significantly associated with the decline of opportunistic parasites and diarrhea ($p < 0.05$) while no ART had relatively greater incidence of opportunistic intestinal parasites.

INTRODUCTION

Gastrointestinal infections are very common in patients with HIV infection or AIDS. Diarrhoea is a common clinical presentation of these infections. Reports indicate that diarrhoea occurs in 30-60 per cent of AIDS patients in developed countries and in about 90 per cent of AIDS patients in developing countries (Kucerova *et al.*, 2011; Missaye *et al.*, 2013). Co-infections of HIV and opportunistic parasites, including intestinal protozoa and helminths, are of concern in resource-poor settings where the health status of the population is generally poor and these opportunistic parasites very common (Tzipori *et al.*, 2008). The resultant effect of such parasitic infections include chronic diarrhea, weight loss, and malnutrition, which has been associated with death among AIDS patients (Tian *et al.*, 2013). HIV has the capacity to circumvent and weaken human immune system providing the impetus for increased infection with parasites such as *Cryptosporidium* spp., *Microsporidium* spp, *Giardia intestinalis*, and *Strongyloides* (S) *stercoralis* (Tian *et al.*, 2013). *Cryptosporidium* spp, *Microsporidia* and other coccidian parasites have emerged as significant causes of persistent diarrhea in People living with HIV/AIDS (PLWHA) (Tumwine *et al.*, 2005). These pathogens have been recognized as worldwide causes of diarrhea in all age groups, yet their most significant impact have been felt among individuals with weakened immune systems, especially PLWHA and organ transplant recipients (Hunter, 2002).

In immunocompromised individuals, diarrheal infections goes beyond the inconvenience of frequent watery stool but may result in severe and potentially life-threatening dehydration, electrolyte loss and malnutrition, and eventually death (Jha *et al.*, 2013). Transmission of *Cryptosporidium* is mainly through the fecal-oral route in contaminated water and food, as well as through person-to-person spread and contact with infected animals (Shimelis *et al.*, 2016). Microsporidiosis, caused by *Microsporidia*, another important opportunistic pathogen causing significant morbidity in PLWHA (Didier *et al.*, 2011). The route of transmission is usually by ingestion of the spores, including evidence of spore inhalation or rectal transmission (10). Antiretroviral therapy increases the length and quality of life and productivity of patients by improving survival and decreasing the incidence of opportunistic infections in PLWHA through the reduction of circulating viremia and increasing the level of CD4+ cells (11). Previous studies in India have investigated intestinal parasitic infections in relation to ART and CD4+ count (Wanyiri *et al.*, 2013; Kulkarni *et al.*, 2009).

MATERIAL AND METHODS

Study patients were interviewed using the structured questionnaire and information was obtained on demographic characteristics, present and past history of diarrhoea and antibiotic treatment.

Diarrhoea was defined as two or more liquid or three or more soft stools per day. Patients already on antibiotic treatment were excluded from the study. A total of 165 patients were enrolled in the study. Blood samples (plain and EDTA) 5 ml each were obtained from enrolled patients. Serum samples were used for HIV testing. HIV serostatus of the patients was determined by using commercially available ELIS Antibody tests (Genetic system, Biorad Labs, USA and Tridot, Mitra and Co., New Delhi) using National AIDS Control Organisation (NACO) recommended algorithm. Patients were categorized by their immune status according to the 1993 – revised classification system for the HIV infection by CD4 T-cell categories (National, 2007; Castro, 1993). Stool specimens were collected according to the WHO standard procedure and examined microscopically following direct and formalin-ether concentration methods (World Health Organization, 1991). Stool samples were collected at home in labelled, leak proof, clean sterile plastic containers and were transported to the laboratory within three hours of collection. The stool samples were fixed in 10 percent formalin saline, concentrated using formyl / ethyl acetate and examined through direct observation insaline (0.85% NaCl solution). Lugol's iodine was used for the detection of ova, larvae, trophozoites and cysts of intestinal parasites. Smears of direct and concentrated specimens were examined by modified acid faststaining for *C. parvum*, *I. belli* and *Cyclospora* (World Health Organization, 1991; Centers for Disease Control and Prevention). Modified trichrome stain (Hi-media laboratories, India, Qualigens Fine Chemicals, India) and modified ZN was used for detecting *Microsporidia* (Weber, 1992) and *Cryptosporidium* (Fig 1) *Statistical analysis*: Data were analysed using SPSS software version 16.0 (SPSS Inc, USA). The proportion of opportunistic pathogens were compared between the ART groups by using Z test.

RESULTS AND DISCUSSION

Among the total of 234 HIV study subjects, 146 (58.2%) males and 94 (41.8%) females were enrolled in the study. Diarrhoea was seen in 142 (60.8%) while 92 (38.2%) (Table 1) were HIV non-diarrhoea as a control group. Overall, 69 cases yielded

parasites 29.4% of which *Cryptosporidium* (21.8%) was the most frequently encountered pathogen in the study population followed by *S.stercoralis* (17.6%), *Ascaris ova* (11%), hookworm (6.6%), trichuris ova (4.8%) (TABLE 2). HIV infected patients, due to down regulation of the immune system. Gastrointestinal parasitic infection is a universally recognized problem in these patients. These infections largely present with diarrhoea leading to life threatening complications. Various studies from India and other countries have reported a high prevalence of intestinal parasite, ranging from 30 to 60 per cent (10,14,21-23). Among the 106 patients with CD4 count < 200 cells / μ l, parasites could be identified in 71 (46%) patients and opportunistic parasites were detected in as many as 69 (41%) patients. Thus, like many other studies, we also found that infections with opportunistic pathogens were the leading cause of diarrhoea in HIV infected individuals, especially, in subjects with advanced disease. *C.parvum* and *I. belli*, were the most common pathogens. Among the non opportunistic pathogens *ascaris*, *hookworm* and *giardia* seed to contribute significantly has shown earlier (Joshi, 2002). Similar to other reports, *Microsporidia* and *Isospora* were detected in a few patients only. There was some limitations in our study.

The study was done on a small sample size as a majority of the patients who came to microbiology laboratory were referred from the general practitioners or from primary or secondary care centers. Patients were also referred from Voluntary Counseling and HIV Testing centers. Majority of the patients seen at these centers had already received antibiotics prior to their visit and therefore the number of symptomatic patients was less. The result is almost similar to the findings from other countries but such differing prevalence rates might be due to differences in geographical location, sensitivity of diagnostic techniques, immune status of study participants, environmental hygiene and possible increased awareness, amongst others (Weber, 1992; Brink *et al.*, 2000). Several species of intestinal opportunistic parasites have been reported among PLWHA (National AIDS Control Organization, 2007), with *Cryptosporidium* spp, *Microsporidium* spp and *S. stercoralis* the most commonly encountered in other study.

Table 1.

Microorganism	ART STATUS		
	NO ART	ON ART	P-VALUE
Cryptosporium(n=19)	13(80.5%)	6(19.5%)	0.001
S.stercoralis(n=16)	11(79.3%)	5(20.7%)	0.006
Ascaris(n=8)	6(66.7%)	2(33.4%)	0.56
Isospora(n=7)	5(71.9%)	2(38.1%)	0.53
Giardia(n=14)	10(71.4%)	4(29.6%)	0.68
Microsporidia spp.(n=3)	3(100%)	0	0.042
Hookworm(n=5)	3(60%)	2(40%)	0.67
Trichuris (n=2)	2(100%)	0	1

Table 2.

Microorganism	Diarrhoeal stool	Non-diarrhoeal stool	p-value
Cryptosporium (n=19)	16(83.3%)	3(17.7%)	0.005
S.stercoralis (n=16)	13(75.8%)	3(24.2%)	0.679
Ascaris (n=8)	6(72.2%)	2(17.8%)	0.221
Isospora (n=7)	6(92.3%)	1(7.7%)	0.003
Giardia (n=14)	9(64.2%)	5(36.8%)	0.833
Microsporidia spp.(n=3)	3(100%)	0	1
Hookworm(n=5)	4(80%)	3(20%)	0.342
Trichuris (n=2)	1(50%)	1(50%)	0.27




Fig 1. Cryptosporidium shown in Modified ZN staining In 1000x

Cryptosporidium spp, which was found to be responsible for diarrhea in 10-20% of PLWHA worldwide (Weber *et al.*, 1999), has been reported in our study, and more importantly in those with CD4+ T-cell counts of less than 200 cells/mm³ (World Health Organization, 1991; Brink *et al.*, 2000).

Conclusion

This study aims to highlight the parasitic causes of diarrhoea among HIV patients. Though it is already known about the causative organisms but it was not known in our area. The impact of ART therapy on the parasitic load is significant in cases of Cryptosporidia and *S.stercoralis*. Therefore, our study emphasizes the need for routine screening of enteric coccidiosis as well as education about practicing personal hygiene and taking timely and appropriate measures. In a developing country like India, the magnitude of intestinal parasitic infections in HIV patients further adds to the existing financial burden of the disease. Patients usually belong to poor socio-economic backgrounds and they can hardly afford treatment. Therefore, it is suggested that steps should be taken to prevent the occurrence of these diseases in AIDS patients, as often the disease may take a fulminant form. This can be done by drinking safe water and avoiding contact with contaminated soil.

REFERENCES

- Brink AK., Mahe C., Watera C., Lugada E., Gilks C., Whitworth, J. *et al.*, 2000. Diarrhea, CD4 counts and enteric infections in a community-based cohort of HIV-infected adults in Uganda. *J Infect* 45 : 99-106.
- Castro KG., Ward JW., Slutsker L., Buehler JW., Jaffe HW., Ruth L. *et al.*, 1993. Revised classification system for HIV infection and expanded surveillance on definition of AIDS among adolescents and adults. *Morbidity and Mortality Weekly Report*, 41: 1-1918. World Health Organization. *Basic laboratory methods in medical parasitology*. Geneva: World Health Organization;1991. p. 9-31.
- Centers for Disease Control and Prevention. DPDx: Laboratory Chacin-Bonilla L., Guanipa N., Cano G., Raleigh X., Quijada L. 1992. Cryptosporidiosis among patients with acquired immunodeficiency syndrome in Zulia State, Venezuela. *Am J Trop Med Hyg.* 47(5): 582-586. PubMed | Google Scholar
- Didier ES., Weiss LM. 2011. Microsporidiosis: not just in AIDS patients. *Curr Opin Infect Dis.* 24(5): 490-495. PubMed | Google Scholar
- Hunter PR., Nichols G. 2002. Epidemiology and Clinical Features of Cryptosporidium Infection in Immunocompromised Patients. *Clin Microbiol Rev.* 15(1): 145-154. PubMed | Google Scholar
- identification of parasites of public concern. Modified 04/06/2001 DPDx; available at www.dpd.cdc.gov/dpdx.
- Jha AK., Uppal B., Chadha S., Aggarwal P., Ghosh R., Dewan R. 2013. Enteric pathogens, immune status and therapeutic response in diarrhea in HIV/AIDS adult subjects from north India. *Curr HIV Res.* 11(4): 326-332. PubMed | Google Scholar
- Joshi M., Chowdhary AS., Dalal PJ., Maniar JK. 2002. Parasitic diarrhea in patients with AIDS. *Natl Med J India.* 15: 72-4.
- Kucerova Z., Sokolova OI., Demyanov AV., Kvac M., Sak B., Kvetonova D. *et al.*, 2011. Microsporidiosis and Cryptosporidiosis in HIV/AIDS Patients in St Petersburg, Russia: Serological identification of microsporidia and Cryptosporidium parvum in sera samples from HIV/AIDS patients. *AIDS Res Hum Retroviruses.* 27(1): 13-15. PubMed | Google Scholar
- Kulkarni SV., Kairon R., Sane SS., Padmawar PS., Kale VA., Thakar MR. *et al.*, 2009. Opportunistic parasitic infections in HIV/AIDS patients presenting with diarrhoea by the level of immunosuppression. *Indian J Med Res.* 130(1): 63-67. PubMed | Google Scholar
- Missaye A., Dagne M., Alemu A., Alemu A. 2013. Prevalence of intestinal parasites and associated risk factors among HIV/AIDS patients with pre-ART and on-ART attending Dessie hospital ART clinic, Northeast Ethiopia. *AIDS Res Ther.* 10(1): 7. PubMed | Google Scholar
- National AIDS Control Organization (NACO). *Manual on quality standards for HIV testing laboratories*. New Delhi: NACO; 2007.
- Ojuromi OT., Izquierdo F., Fenoy S., Fagbenro-Beyioku A., Oyibo W., Akanmu A. *et al.*, 2012. Identification and characterization of microsporidia from fecal samples of HIV-positive patients from Lagos, Nigeria. *PloS One.* 7(4): e35239. PubMed | Google Scholar
- Ojuronbe O., Raji OA., Akindele AA., Kareem MI., Adefioye OA., Adeyeba AO. 2011. Cryptosporidium and other enteric parasitic infections in HIV-seropositive individuals with and without diarrhoea in Osogbo, Nigeria. *Br J Biomed Sci.* 68(2): 75-78. PubMed | Google Scholar
- Satheesh Kumar, S., Ananthan S., Lakshmi P. 2002. Intestinal parasitic infection in HIV infected patients with diarrhea in Chennai. *Indian J Med Microbiol.* 20 : 88-91.
- Shimelis T., Tassachew Y., Lambiyu T. 2016. Cryptosporidium and other intestinal parasitic infections among HIV patients in southern Ethiopia: significance of improved HIV-related care. *Parasit Vectors.* 9: 270. PubMed | Google Scholar
- Stentiford GD., Becnel JJ., Weiss LM., Keeling PJ., Didier ES., Williams BAP. *et al.*, 2016. Microsporidia - Emergent Pathogens in the Global Food Chain. *Trends Parasitol.* 32(4): 336-348. PubMed | Google Scholar
- Tian LG., Wang TP., Lv S., Wang FF., Guo J., Yin XM. *et al.*, 2013. HIV and intestinal parasite co-infections among a Chinese population: an immunological profile. *Infect Dis Poverty.* 2(1): 18. PubMed | Google Scholar
- Tumwine JK., Kekitiinwa A., Bakeera-Kitaka S., Ndezi G., Downing R., Feng X. *et al.*, 2005. Cryptosporidiosis and microsporidiosis in Ugandan children with persistent diarrhea with and without concurrent infection with the

- human immunodeficiency virus. *Am J Trop Med Hyg.* 2005; 73(5): 921-92. PubMed | Google Scholar
- Tzipori S., Widmer G. 2008. A hundred-year retrospective on cryptosporidiosis. *Trends Parasitol.* 24(4): 184-189. PubMed | Google Scholar
- Wanyiri JW., Kanyi H., Maina S., Wang DE., Ngugi P., O'Connor R. *et al.*, 2013. Infectious diarrhoea in antiretroviral therapy-naive HIV/AIDS patients in Kenya. *Trans R Soc Trop Med Hyg.*, 107(10): 631-638. PubMed | Google Scholar
- Weber R., Bryan RT., Owen RL., Wilcox CM., Gorelkin L., Visvesvara GS. 1992. Improved light- microsporidial detection of *Microsporidia* spores in stool and duodenal aspirate. The Enteric Opportunistic Infections Working Group. *N Engl J Med* 326 : 161-6.
- Weber R., Ledergerber B., Zbindin R., Altwegg M., Pfyffer GE., Spycher MA. *et al.*, 1999. Enteric infections and diarrhea in human immunodeficiency virus- infected persons: prospective community-based cohort study. Swiss HIV cohort study. *Arch Intern Med* 159 : 1473-80.
- Wiwanitkit V. 2001. Intestinal parasitic infections in Thai HIV infected patients with different immunity status. *BMC Gastroenterol.*, 1 : 3-5.
